



Wash the face with lukewarm water



Apply anesthetic cream on injection area

**PREPARATION
FOR
INJECTION**




Leave it for 20-30 minutes



Wipe off the cream with an alcohol cotton pad


1 syringe



Eye Protocol

● x2 / 0.5cc per point


1+1 syringe



GENERAL Protocol

● x2 / 0.2cc per point
● x8 / 0.2cc per point

1+1+1 syringe



Enhancing Protocol


● x2 / 0.5cc per point
● x8 / 0.25cc per point

HOW TO INJECT GOURI

General Protocol

- ⦿ Inject to deep dermis
- ⦿ Inject 0.2 cc per point


- 1st point: Middle point of the forehead, above the center of an eye pupil
- 2nd point: Horizontal with nostrils, about 2-3 cm away from the nose on the medial cheek
- 3rd point: Same level as 2nd point, below the cheekbone
- 4th point: Same level with the end of the lips line on the superior jawl,
located in the middle between the 2nd and 3rd point
- 5th point: Inject one point on the red marked areas,
about 1.5~2.0cm horizontally and vertically away from the of the eyes.



HOW TO INJECT GOURI

Eye Protocol

- ☐ Inject to deep dermis
- ☐ Inject 0.5 cc per point



- Injection point : Inject one point on the red marked areas, about 1.5~2.0cm horizontally and vertically away from the of the eyes.

HOW TO INJECT GOURI



Enhancing Protocol

☉ Inject to deep dermis

☉ Inject 0.5 cc per point - eyes

Inject 0.25 cc per point – entire face

- If you want special care, inject the same point of General protocol with 0.25cc for 1st ~ 4th point and with 0.5cc for eyes



PRECAUTIONS FOR USE

- Avoid exposure to heat and radiation (sauna, sun), swimming/bathing and alcohol consumption for the first 24 hours.
- An anesthetic cream or local anesthetic injection can be given before the procedure at the discretion of the doctor.
- Avoid Blood Vessels while Injecting and take special care when you treat under eyes.
- If you have a patient with allergy or Irritable Syndrome, we recommend taking Cetirizine (Prescription-Free) 30 minutes to 1 hour before treatment.
- Not recommended to mix with other products.

PRECAUTIONS RELATED TO COVID 19

- Avoid treatment for 1 month before and after vaccination.
- Cetirizine is recommended 30 minutes to 1 hour before treatment.

STORAGE AND AFTER USE

- Store at room temperature up to 30°C and recommend to use it all on the same day if possible after opening.
- After use, make sure to dispose of it.